## Children 7 – 10 Years of Age Need These Immunizations to Attend School in Arizona 2008-2009 School Year 4th Dose 2<sup>nd</sup> Dose 3rd Dose 5<sup>th</sup> Dose **Immunizations** 1<sup>st</sup> Dose (DTaP/DTP) Diphtheria, Tetanus & Pertussis, Required if dose #1 was given (DT) Diphtheria & Required Required Required before 12 months of age. Tetanus, Td (Tetanus & Diphtheria An interval of at least 4 weeks is required between the first doses of DTaP/DT/Td. An interval of 6 months is required between the last two doses. (IPV/OPV) Polio Not needed if dose #3 was given at 4+years. Required Required Required 4 doses meet the requirement even if all were given in the first year of life. An interval of at least 4 weeks is required between all **Polio** doses. (MMR) Measles, Mumps, & Rubella Required Required MMR #1 is due at 12-15 months of age. MMR #2 may be given 1 month after MMR #1. MMR #2 is required for grades K-12. (Hep B) Hepatitis B Required. Required only if dose #3 was Required Required Must be at least 24 weeks given before 24 weeks of age. of age. Hep B #1 may be given at birth. Dose #2 is due 4-8 weeks after dose #1. Dose #3 is due 2-5 months after dose #2, at least 4 months after dose #1 and no sooner than 24 weeks of age. A 2-dose adolescent series may be given to children 11-15 years of age over a 4-6 month period. Varicella (Chicken Pox) Required Recommended In the 2008-2009 school year, children enrolled in preschool, kindergarten and grades 1, 2, 3, 4, 7, 8, 9, and 10 are required to be immunized against varicella (chicken pox) or have a history of chicken pox disease. Parental recall of chicken pox is acceptable. One dose of varicella vaccine meets the requirement for all children except those vaccinated at 13+ years of age. Note: If MMR and Varicella vaccine doses were not given on the same day, they must be given at least 28 days apart.